

SEPTEMBER 19, 2012

Multi-fractional spacetimes, asymptotic safety and Hořava–Lifshitz gravity

Gianluca Calcagni

Instituto de Estructura de la Materia, IEM-CSIC, Serrano 121, 28006 Madrid, Spain

E-mail: calcagni@iem.cfmac.csic.es

ABSTRACT: We compare the recently formulated multi-fractional spacetimes with field theories of quantum gravity based on the renormalization group (RG), such as asymptotic safety and Hořava–Lifshitz gravity. The change of spacetime dimensionality with the probed scale is realized in both cases by an adaptation of the measurement tools (‘rods’) to the scale, but in different ways. In the multi-fractional case, by an adaptation of the position-space measure, roughly corresponding to a scale dependence of the coordinates. In the case of RG-based theories, by an adaptation of the momenta. The two pictures are mapped into each other, thus presenting the fractal structure of spacetime in RG-based theories under an alternative perspective.

KEYWORDS: Models of Quantum Gravity, Renormalization Group, Fractal Geometry

Contents

1	Introduction	1
2	Three approaches to quantum geometry	4
2.1	Multi-fractional spacetimes	4
2.2	Asymptotically-safe quantum gravity	6
2.3	Hořava–Lifshitz gravity	7
3	Multi-fractional picture of the RG flow	8
3.1	Multi-fractional RG flow and asymptotic safety	8
3.2	Multi-fractional RG flow and Hořava–Lifshitz gravity	12
4	Discussion	13

1 Introduction

When geometry becomes quantum, the dimension of effective spacetime changes with the scale. This phenomenon has been dubbed dimensional reduction [1–3] or dimensional flow [4], and it appears across a variety of approaches both in the continuum (such as in non-commutative spacetimes, asymptotic safety, Hořava–Lifshitz gravity and fractal field theory) and in discrete settings (causal dynamical triangulations, spin foams and loop quantum gravity among others). While in the latter category of models it is often problematic to get the continuum limit of a discrete geometry, the former class can be manipulated more easily and it is particularly suitable to make contact with physics and extract some phenomenology across different scales.

Multi-scale phenomena are not a foreign topic in quantum field theory thanks to the massive deployment of renormalization group (RG) techniques in the most various contexts. When geometry itself is quantized, however, dimensional reduction acquires certain universal traits and both its underlying mechanism and the precise relation with the RG flow need to be explored thoroughly. This problem can be naturally studied in fractal field theory [4–6], especially in its fractional incarnation [7–13] (see [14] for a review). In this case, one can easily tune the ingredients of a Lebesgue–Stieltjes fractal field theory of fixed dimensionality living in fractal Minkowski spacetime $\mathcal{M}_v^D$, the latter being embedded in D -dimensional ordinary Minkowski spacetime M^D . Namely, these ingredients are: (i) the measure weight $v(x)$ in position space and, from that, the Hausdorff dimension $d_H(\mathcal{M}_v^D)$ of position space [8, 9]; (ii) the measure weight $w(p)$ in momentum space and, from that, the Hausdorff dimension $d_H(\mathcal{M}_w^D)$ of momentum space [10, 12]; (iii) the Laplacian operator $\mathcal{K}$ in position space [9, 10, 12]; (iv) the type of diffusion (normal or anomalous) associated with $\mathcal{M}_v^D$ [12]. From (i), (iii) and (iv), one can extract (v) the spectral dimension $d_S(\mathcal{M}_v^D)$

of spacetime [8, 12]. This geometry can be extended to a multi-scale setting [9, 12] and fields can be introduced thereon. Dimensional flow is fully analyzable and one can exercise quantitative control at all scales, not just in isolated regimes along the flow [9, 11]. Also, both at different points in the flow and as a whole, diffusion in quantum spacetime is given a physical and probabilistic interpretation in terms of stochastic processes [12].

Although fractal spacetimes and field theories have been proposed as fundamental, an alternative point of view is to regard them as *effective models* describing certain (regimes of) other, independent quantum-spacetime/quantum-gravity theories. The strategy is to identify the above five data in the fundamental theory we are interested in. In the known cases, we do not have all of them at hand. In theories defined on group manifolds (e.g., non-commutative spaces), an additional complication is to extract analogs of (i)–(v).¹ For theories living in a continuum, such as asymptotic safety (also known as quantum Einstein gravity, QEG [16, 17]; see [18–21] for reviews), this difficulty is partially removed but some of the data may be missing or undecided.² Next, we match these data with the corresponding ones of the fractional theory. Even when both models are on the continuum and on $\mathbb{C}$ -numbers, this step is not always just a matter of recasting the description of the fundamental theory in terms of the language of fractal geometry. Rather, fractal geometry can give constructive insight to fill the gaps in the fundamental theory, remove ambiguities in previous guesswork, and clarify the physical meaning of the ingredients. Last, one extends the mapping to multi-fractal geometries where the effective spacetime dimension varies with the scale.

Due to the similarity of dimensional flow in QEG, multi-fractional spacetimes and Hořava–Lifshitz field-theory approach [23, 24], one may ask whether the multi-scale (in particular, multi-fractal) geometry of multi-fractional spacetimes can serve as an alternative or complementary description of dimensional flow in these and other RG-based theories. The answer is in the affirmative. In the present paper, we gain some insight on the nature of the RG flow by interpreting a multi-scale geometry as a continuous hierarchical adaptation of ‘rods’ (i.e., geometry measurements) on the length scale of the probed phenomena, with respect to the scale of classical macroscopic rods. Thus, dimensional reduction stemming from RG techniques admits a ‘dual’ description in terms of multi-fractal geometry.

It is important to stress that the scope of this dual interpretation is limited by the physical assumptions, i.e., by the symmetries of the system. In particular, from the symmetry choice different Laplacians lead to physically inequivalent models. We take the three examples of multi-fractional spacetimes, QEG and Hořava–Lifshitz gravity for illustration, in the absence of curvature (the metric is the Minkowski one, $g_{\mu\nu} = \eta_{\mu\nu} = \text{diag}(-, +, \dots, +)$). In all three cases, the Laplace–Beltrami operator $\mathcal{K}$ converges in the infrared (IR) to the

¹A wide class of non-commutative spacetimes can be mapped to fractional geometries by recognizing that the classical cyclicity-invariant measure in position space, associated with a given (possibly non-linear) spacetime algebra, is of fractional type [15]. In particular, this helped in reinterpreting the problematic cyclic measure $d\mathbf{x}/(\prod_i x_i)$ of κ -Minkowski spacetime as the asymptotic limit of a more complicated geometry.

²In the case of QEG, (i) is known at any scale, (ii) is assumed, (iii) is tailored compatibly with the RG flow, (v) is known at the UV and IR fixed points, and (iv) can be inferred consequently [22].

usual d'Alembertian,

$$\mathcal{K} \stackrel{\text{IR}}{\sim} \square = -\partial_t^2 + \nabla^2, \quad (1.1)$$

where ∇^2 is the spatial Laplacian. At scales between the IR and the ultraviolet (UV), we have the following limits.

- *Multi-fractional spacetimes.* To get a two-dimensional geometry in the UV [9], one can choose the second-order self-adjoint operator

$$\mathcal{K}_\alpha = \eta^{\mu\nu} \mathcal{D}_\mu \mathcal{D}_\nu, \quad \mathcal{D}_\mu := \frac{1}{\sqrt{v_\alpha(x)}} \partial_\mu \left[\sqrt{v_\alpha(x)} \cdot \right], \quad (1.2)$$

where $v_\alpha(x)$ is the spacetime measure weight at a given scale governed by the parameter α . In the UV, $\alpha = 2/D$, where D is the number of topological dimensions. Poincaré invariance is broken except in free models [25]. Renormalization can be studied in a perturbative quantum field theory context.

- *Asymptotic safety.* Lorentz invariance is preserved throughout the RG/dimensional flow and, at any given scale, the Laplacian is effectively given by

$$\mathcal{K}_\delta \sim \square^{1+\frac{\delta}{2}}, \quad (1.3)$$

where $\delta = 2$ at the UV non-Gaussian fixed point, $\delta = 0$ in the IR and it can acquire other non-trivial values in between. Equation (1.3) is meant to be only semi-quantitative, since the actual scaling takes place at the level of the metric and in momentum space. The theory is studied non-perturbatively via the functional renormalization group approach.

- *Hořava–Lifshitz spacetimes.* Here Lorentz invariance is preserved only on spatial slices and time is a preferred direction. The Laplace–Beltrami operator is then

$$\mathcal{K}_z = -\partial_t^2 + \nabla^{2z}, \quad (1.4)$$

where z is a parameter fixed at $z = D - 1$ in the UV and, effectively, $z = 1$ in the IR, eq. (1.1). As in multi-fractional models, renormalization is formulated perturbatively.

Even if these models are characterized by a similar or identical dimensional flow, they are essentially different in other predictions. However, the Laplacian operator does not enter the geometric description presented in this paper, which is rather general. Therefore, the latter may be regarded not as a physical duality between theories, but as a complementary geometric description of their multi-scale properties related to momentum space.

In section 2 we briefly review multi-fractional, QEG and Hořava–Lifshitz spacetimes. Section 3 is the core of the paper. We shall concentrate on the comparison between multi-fractional theories and asymptotic safety; the case of Hořava–Lifshitz stems from a slight generalization of the obtained results. A discussion follows in section 4.

2 Three approaches to quantum geometry

2.1 Multi-fractional spacetimes

Without loss of generality we concentrate on fractional Minkowski spacetime $\mathcal{M}_v^D = \mathcal{M}_\alpha^D$, where the metric is $\eta_{\mu\nu}$ and there is no gravity. $\mathcal{M}_\alpha^D$ lives in D -dimensional ordinary Minkowski spacetime M^D , (spanned by coordinates x^μ , $\mu = 0, 1, \dots, D-1$) but, contrary to the latter, it is endowed with a non-trivial measure of the form

$$d\varrho_\alpha(x) = d^D x v_\alpha(x), \quad (2.1a)$$

$$v_\alpha(x) = \prod_\mu v_\alpha(x^\mu) \quad (2.1b)$$

$$:= \prod_\mu \frac{|x^\mu|^{\alpha_\mu - 1}}{\Gamma(\alpha_\mu)}, \quad (2.1c)$$

where Γ is the gamma function and α_μ are D real parameters (‘fractional charges’) in the (further restrictable) range $0 < \alpha_\mu \leq 1$. From the scaling law of the measure, $\varrho_\alpha(\lambda x) = \lambda^{d_H} \varrho_\alpha(x)$, one infers that the Hausdorff dimension of fractional spacetime is $d_H = \sum_\mu \alpha_\mu$, as one can show also with other independent calculations [8]. The simplest type of fractional measure is ‘isotropic’ and $\alpha_\mu = \alpha$ are all equal; we shall focus our attention on this configuration except when discussing Hořava–Lifshitz spacetimes. It is also possible to extend the geometric construction to complex-valued fractional charges [7, 9, 15] or more general factorizable measures [13, 25] but we shall not do so since scaling properties are more apparent in the real-order fractional case.

The measure can be reduced to the Lebesgue one by defining a set of coordinates, which we call ‘geometric,’ such that

$$q^\mu := \varrho_\alpha(x^\mu) = \frac{\text{sgn}(x^\mu) |x^\mu|^\alpha}{\Gamma(\alpha + 1)}, \quad (2.2)$$

so that $d\varrho_\alpha = d^D q$. Geometric coordinates will be essential to compare fractional theories with other proposals.

Fractional spacetimes can be equipped with generalizations of the Laplace–Beltrami operator (1.1). A particular choice is the second-order operator (1.2), which is self-adjoint with respect to the natural scalar product [10]. Laplacian operators with fractional order $2\gamma \neq 2$ [12] or second-order but with different measure factors [9, 10] are also possible.

Momentum space $\mathcal{M}_w^D = \mathcal{M}_{\alpha'}^D$ may be equipped with a different fractional measure $w(p_{\text{frac}}) = v_{\alpha'}(p_{\text{frac}})$. In the following we dub as p_{frac} the physical momentum in fractional models. The functions

$$\mathfrak{e}(p_{\text{frac}}, x) = \frac{1}{\sqrt{v_{\alpha'}(p_{\text{frac}}) v_\alpha(x)}} \frac{e^{ip_{\text{frac}} \cdot x}}{(2\pi)^{\frac{D}{2}}} \quad (2.3)$$

are eigenfunctions of (1.2), $\mathcal{K}_\alpha \mathfrak{e}(p_{\text{frac}}, x) = -p_{\text{frac}}^2 \mathfrak{e}(p_{\text{frac}}, x)$, where $p_{\text{frac}}^2 := \eta_{\mu\nu} p_{\text{frac}}^\mu p_{\text{frac}}^\nu = -(p_{\text{frac}}^0)^2 + \sum_{i=1}^{D-1} (p_{\text{frac}}^i)^2$. They form a basis for the unitary invertible momentum transform

$$\tilde{f}(p_{\text{frac}}) := \int_{-\infty}^{+\infty} d\varrho_{\alpha}(x) f(x) \mathfrak{e}^*(p_{\text{frac}}, x) =: F_{\varrho}[f(x)], \quad (2.4a)$$

$$f(x) = \int_{-\infty}^{+\infty} d\varrho_{\alpha'}(p_{\text{frac}}) \tilde{f}(p_{\text{frac}}) \mathfrak{e}(p_{\text{frac}}, x). \quad (2.4b)$$

If momentum and position space have the same measure ($\alpha = \alpha'$), F_{ϱ} is an automorphism. The identity resolution of this transform is the fractional generalization of the delta distribution, $\delta_{\alpha}(x, x') := \delta(x - x') / \sqrt{v_{\alpha}(x)v_{\alpha}(x')} = \int d\varrho_{\alpha'}(p_{\text{frac}}) \mathfrak{e}^*(p_{\text{frac}}, x) \mathfrak{e}(p_{\text{frac}}, x')$.

Dynamics enters when an action is defined. For example, a scalar field theory is given by $S_{\alpha} = \int d\varrho_{\alpha}(x) [(1/2)\phi \mathcal{K}_{\alpha} \phi - V(\phi)]$ for some potential V . The Laplacian also determines the spectral dimension d_S of spacetime, defined as the trace in position space of the heat kernel operator. It represents the dimensionality felt by a test particle left to diffuse for an infinitesimal time. For normal (i.e., non-fractional) diffusion, one can show that $d_S = d_H$ [8]. If the diffusion operator is fractional with order $0 < \beta \leq 1$, then the diffusion equation is of the form $(\partial_{\sigma}^{\beta} - \mathcal{K}_{\alpha})P = 0$ (where σ is a fictitious diffusion time or scale and P is the probability distribution governing the process) and $d_S = \beta d_H$ [12].

In order to obtain a multi-scale geometry, it is sufficient to replace the measure ϱ_{α} with a linear superposition of fractional measures, summed over a finite number of parameters $0 < \alpha_n \leq 1$ [7, 9, 12]:

$$\varrho(x) = \sum_{n=1}^N g_n \varrho_{\alpha_n}(x), \quad (2.5)$$

where g_n are some dimensionful coupling constants containing the scales of the system. This type of multi-component measures is encountered in multi-fractal geometry and complex systems (see [26] and [9, 12] for further details). The action becomes a sum of α -dependent Lagrangians, $S = \sum_n g_n S_{\alpha_n}$, while the diffusion equation is realized by summing over α and also β [11, 12]. Fixing β for simplicity, one has $(\partial_{\sigma}^{\beta} - \sum_{n=1}^N \zeta_n \mathcal{K}_{\alpha_n})P = 0$, where the coefficients ζ_n have the form $\zeta_1(\ell) = (\ell_1/\ell)^2$, $\zeta_n(\ell) = [\ell_n/(\ell - \ell_{n-1})]^2$, $\zeta_N \equiv 1$ [11, 12]. Here $\ell = \ell_N$ is the largest scale, which is identified with the probed scale in a measurement. The spectral dimension is approximately given by $d_S(\ell) \approx \beta D \alpha_{\text{eff}}(\ell)$, where the effective fractional charge reads

$$\alpha_{\text{eff}}(\ell) = \frac{1 + \sum_{n=1}^{N-1} \zeta_n(\ell) \alpha_n}{1 + \sum_{n=1}^{N-1} \zeta_n(\ell)}. \quad (2.6)$$

The most important examples are also the simplest ones. When $N = 2$, the spectral dimension flows monotonically between two limiting values $d_S \sim D\alpha_1$ and $d_S \sim D\alpha_2$. Setting $D = 4$, $\alpha_1 = 1/2$ and $\alpha_2 = 1$, we get a flow from 2 to 4 dimensions. When $N = 3$, there are two scales and three values of α_n which reproduce (for $D = 4$, $\alpha_1 = 1/2$, $\alpha_2 = 1/3$, $\alpha_3 = 1$) the asymptotic regimes of QEG in the presence of a cosmological constant [11, 12, 22].

The polynomial measure (2.5) guarantees that geometry changes with the scale. As a side remark, we notice that it automatically embodies the idea that, in a multi-fractal

spacetime, coordinates acquire a scale dependence [27, 28]. Using eq. (2.6), one can in fact identify an effective geometric coordinate system (index μ omitted)

$$q_{\text{eff}}(\ell) \approx \frac{\text{sgn}(x)|x|^{\alpha_{\text{eff}}(\ell)}}{\Gamma[\alpha_{\text{eff}}(\ell) + 1]} \sim \text{sgn}(x)|x|^{\frac{d_H(\ell)}{D}}, \quad (2.7)$$

which replaces eq. (2.2).

Multi-fractional spacetimes are models of quantum geometry in two distinct ways. As a fundamental theory, by prescribing a certain action and then quantizing gravity perturbatively on the same footing than other fields. As a phenomenological model, by regarding these spacetimes as effective descriptions of geometry valid in certain regimes of a given fundamental theory. In the second case (which we will implicitly assume in section 3) the details of the action are engineered according to the guidelines of the fundamental theory. As anticipated in the introduction, this tailoring procedure constrains the choice of symmetries and of Laplacian and, ultimately, it boils down to a choice of momentum space.

2.2 Asymptotically-safe quantum gravity

Quantum Einstein gravity aims to quantize the gravitational interaction as a field theory. Although graviton perturbation theory suffers from divergences, the program can be carried out non-perturbatively via the functional renormalization group approach. This method has wide applications (see [29–32] for general reviews) but here we only recall the main results in QEG (e.g., [20, 21] and references therein).

To uniform our notation with the literature, we use the symbol k for the momentum/energy scale representing a cut-off above which all quantum fluctuations are integrated out. For a given k , effective QEG spacetime and dynamics are described by the average action $\Gamma_k[g_{\mu\nu}]$. The metric is subject to quantum fluctuations and, depending on the resolution $L \sim k^{-1}$ of the microscope probing the geometry, one will observe different properties of spacetime. Thus, the system is multi-scale. The change of the microscope amounts to a coarse-graining procedure. Denote as $\langle g_{\mu\nu} \rangle_k$ the metric averaged over Wick-rotated spacetime volumes of linear size $L \sim k^{-1}$. Then, $\langle g_{\mu\nu} \rangle_k$ is solution of

$$\frac{\delta \Gamma_k}{\delta g_{\mu\nu}} [\langle g_{\mu\nu} \rangle_k] = 0. \quad (2.8)$$

Letting k take any non-negative value, one ends up with a continuous family of actions and of field equations (2.8), all valid simultaneously. From the scale dependence of Γ_k , one can extract the solution $\langle g_{\mu\nu} \rangle_k$ at any scale from the UV to the IR. Although $\langle g_{\mu\nu} \rangle_k$ is typically a smooth classical metric, its change with the scale, regulated by the renormalization group flow, makes effective QEG spacetime a non-smooth multi-scale (in particular, fractal [20]) object with possibly very ‘irregular’ geometry.

In the absence of matter and in the so-called Einstein–Hilbert truncation of Γ_k , the scale dependence of the average metric can be recast in terms of the running cosmological constant $\bar{\lambda}_k$. In fact, the field equations read $R_{\mu\nu}[\langle g \rangle_k] = [2/(2 - D)]\bar{\lambda}_k \langle g_{\mu\nu} \rangle_k$. Let k_0

be an arbitrary reference scale (typically in the infrared, $k_0/k \gg 1$) and assume that the cosmological constant scales according to a function F ,

$$\bar{\lambda}_k = F(k^2) \bar{\lambda}_{k_0}, \quad (2.9)$$

where F depends on k^2 by the requirement of Lorentz invariance. We also assume that F stays positive throughout the flow between k and k_0 . Then, from $F^{-1} R^\mu{}_\nu[\langle g \rangle_k] = [2/(2-D)] \bar{\lambda}_{k_0} \delta^\mu_\nu$ and the scaling property $R^\mu{}_\nu[cg] = c^{-1} R^\mu{}_\nu[g]$ of the Riemann tensor ($c > 0$ is a constant), one gets

$$\langle g_{\mu\nu} \rangle_k = \frac{1}{F(k^2)} \langle g_{\mu\nu} \rangle_{k_0}, \quad \langle g^{\mu\nu} \rangle_k = F(k^2) \langle g^{\mu\nu} \rangle_{k_0}. \quad (2.10)$$

In particular, from the inverse metric it follows that the Laplacian $\Delta(k) = (\langle g \rangle_k)^{-1/2} \times \partial_\mu (\langle g \rangle_k^{1/2} \langle g^{\mu\nu} \rangle_k \partial_\nu)$ scales as

$$\Delta(k) = F(k^2) \Delta(k_0). \quad (2.11)$$

The function F acquires different asymptotic forms depending on the regime. In the far infrared, $F \sim 1$ by definition ($k \rightarrow k_0 \rightarrow +\infty$). In an intermediate semi-classical regime near the Gaußian fixed point, $F \sim k^4$. In the deep UV, at the non-Gaußian fixed point, asymptotic safety implies that $F \sim k^2$ [33]. All in all, we can write

$$F \sim |k|^\delta \quad (2.12)$$

in asymptotic regimes (plateaux of the spectral dimension $d_S = 2D/(2+\delta)$), with $\delta = 0, 4, 2$ in the IR, semi-classical limit and UV, respectively. The semi-classical limit may change depending on the truncation and on the presence of matter [34]. Outside asymptotic regimes, the details of dimensional flow are non-universal and depend on the regularization scheme [12, 22]. Notice that the effective spacetime measure is the standard Lebesgue measure $d^D x \sqrt{|g|}$, so in the zero-curvature limit the Hausdorff dimension coincides with the topological one, $d_H = D$.

2.3 Hořava–Lifshitz gravity

In Hořava–Lifshitz spacetimes, anomalous scaling is associated with the coordinates rather than the metric. In anisotropic ‘critical’ systems, coordinates scale as

$$t \rightarrow \lambda^z t, \quad \mathbf{x} \rightarrow \lambda \mathbf{x}, \quad (2.13)$$

for constant λ , so that time and space directions have dimensions $[t] = -z$ and $[x^i] = -1$ in momentum units. Anisotropic scaling between time and space appears in systems such as, e.g., Lifshitz scalar field theory [35–39],

$$S_{\text{Lifshitz}} = \frac{1}{2} \int dt d^{D-1} x \left[\dot{\phi}^2 - \frac{1}{4} (\Delta \phi)^2 \right], \quad (2.14)$$

where a dot denotes a derivative with respect to time t and $\Delta = \partial_i \partial^i$ is the spatial Laplacian. In this example, the dynamical critical exponent (or anisotropic scaling exponent) is $z = 2$. Its value characterizes the critical behaviour of correlation functions

of the scalar field near a phase transition, as one can infer from the conformal dimension $[\phi] = (D-1-z)/2$. When $D-1 = z$ (two spatial dimensions), the field propagator becomes logarithmic and the system is said to be at quantum criticality. Examples of multi-critical systems are certain metamagnets, liquid crystals and Ising models. These can be studied with RG techniques; the Lifshitz scalar theory (2.14) possesses two Gaussian fixed points, one at $z = 2$ where the system is invariant under the anisotropic scaling (2.13) and one at $z = 1$ where the operator $(\Delta\phi)^2$ becomes irrelevant and local Lorentz invariance can be restored by a relevant operator $\sim \phi\Delta\phi$.

In the case of gravity [23], one constructs a $z = D - 1 = 3$ action with all possible relevant operators respecting foliated diffeomorphisms (i.e., spatial rotations and time reparametrizations). We do not write the action because there exist several inequivalent proposals, depending on the assumptions [40]. In general, there are up to $O(\partial^{2z})$ operators both in the gravitational and matter sector. The spectral dimension turns out to be $d_S = 1 + (D-1)/z$ [24, 41]. Thus, in the UV $d_S = 2$. As the Lifshitz scalar example suggests from the behaviour of the propagator, the advantage in formulating gravity as a critical anisotropic system is that the theory is perturbatively (power-counting) renormalizable. Lorentz invariance is recovered, at least at the classical level (but see [42]), in the infrared.

3 Multi-fractional picture of the RG flow

Multi-fractional theory and renormalization group flow are multi-scale systems with similar characteristics. The purpose of this section is to show that this similarity is not accidental. The idea is simple and is based on the observation that in the RG flow the physical momentum carries the scale dependence.³ In multi-fractional theory, multi-scaling is realized in position space by a scale-dependent measure and, hence, effectively scale-dependent coordinates. In turn, this translates into scale-dependent conjugate momenta, which can be compared with the momenta in other RG-based theories.

3.1 Multi-fractional RG flow and asymptotic safety

A first cursory look at multi-fractional spacetimes often inspires the following crude correspondence with a covariant theory. Comparing multi-fractional actions with any covariant action, the measure weight $v(x)$ in flat space mimicks the determinant factor $\sqrt{-g}$ on a curved background. Any anomalous scaling of the metric $g_{\mu\nu}$ would survive in a local inertial frame, in which case it can be ascribed, effectively, to a non-trivial measure weight.

At the effective-action level, for QEG effective spacetimes we can make the argument more precise (on a local inertial frame). We first develop it for *fixed* dimensionality. As we have seen, in D dimensions the QEG average metric $g_{\mu\nu}(k) = \langle g_{\mu\nu} \rangle_k$ scales as

$$g_{\mu\nu}(k) = |k|^{-\delta} g_{\mu\nu}(k_0), \quad (3.1)$$

where $|k|$ is the norm of the vector k^μ in imaginary time representing the IR cut-off of the theory. Wick-rotating back to Lorentzian signature, the scaling law is presented in a

³For instance, in perturbative RG a change in the momentum with the scale can be seen as a modification of the kinetic term, corresponding to field-strength (wave-function) renormalization.

Lorentz-invariant way. Preservation of Lorentz invariance guarantees control over operator proliferation along the RG flow. Equation (3.1) implies $\sqrt{-g} = |k|^{-D\delta/2} \sqrt{-g_0}$. From now on, let us identify the RG cut-off scale k with the physical momentum,

$$k \equiv p_{\text{QEG}}; \quad (3.2)$$

this minimal identification is employed in the literature and does not result in a loss of generality for our purposes. In fractional theory, the transform (2.4) maps functions in position space to functions in physical momentum space, so that the momentum p_{frac} is conjugate to the coordinate x , $p_{\text{frac}} \leftrightarrow x^{-1}$. If we identified the physical momenta of both theories with each other,

$$p_{\text{QEG}} \stackrel{?}{\sim} p_{\text{frac}} \quad (3.3)$$

for each direction, then $k = p_{\text{QEG}} \leftrightarrow x^{-1}$ and the non-trivial measure in the effective action Γ_k would be $'x^{D\delta/2}'$, where this writing loosely represents the Euclidean modulus $|x|^{D\delta/2}$ or a generic power of the coordinates such as $|x_0 \dots x_{D-1}|^{\delta/2}$. By itself this point is unclear, but further problems arise on the fractional side, where the measure is $|x_0 \dots x_{D-1}|^{\alpha-1}$. This is a negative power, which could never match with $\delta/2 \geq 0$.⁴ To get the correct matching, reproducing the dimensional flow and the associated profile of the spectral dimension, we should set

$$\alpha = \frac{2}{2 + \delta}, \quad (3.4)$$

implying the identification $p_{\text{QEG}} \leftrightarrow x^{-\alpha} \sim q^{-1}$ and, therefore,

$$p_{\text{QEG}} \sim (p_{\text{frac}})^\alpha. \quad (3.5)$$

This relation replaces (3.3) and, *a posteriori*, is more natural than any other. In fact, $\{q\}$ are nothing but the geometric coordinates (2.2) of fractional spaces. In these special coordinates, the measure becomes an ordinary Lebesgue measure $d^D q$, with Poincaré symmetries acting on q (which are, therefore, non-linear transformations on x). This also fixes the mismatch between the Lorentz-invariant QEG framework and the Lorentz-breaking fractional setting. To carry out the fractional dual picture of asymptotic safety, *we must identify QEG spacetime coordinates with fractional geometric coordinates*.

Equation (3.5) states that the physical momentum p_{QEG} in QEG is mapped to the inverse scale conjugate to q . Giving length units to both types of coordinates by associating them with measurement scales,

$$x \leftrightarrow p_{\text{frac}}^{-1} \sim \ell, \quad q \leftrightarrow p_{\text{QEG}}^{-1} \sim L, \quad (3.6)$$

their role is to define the ‘rods’ at a given scale. x has engineering dimension -1 and so ℓ is a scale at which measurements are taken with a classical, macroscopic rod. The scaling of q is different, so L is not a length scale but a power of length, $L \sim \ell^\alpha$. But this is nothing but the ‘length’ of a natural rod at the scale where the dimension is (for each direction) α .

⁴If we compared this measure with the effective measure $'|x|^{-2\delta}'$ of Γ_{k_0} , we would not fare better. In that case, $\alpha = 1 - \delta/2$, which does not do the job since $\alpha > 0$ but $\delta \geq 2$ in the UV and semi-classical regimes.


We can give a pictorial reformulation of this statement, first for a fractal with fixed dimension embedded in an ambient space, and then for a multi-fractal.

- *Fixed dimension.* Imagine an observer $\mathcal{O}_{\mathcal{F}}$ living in a fractal space $\mathcal{F}$ and making measurements with a rod. Call the latter a ‘ q -rod,’ measuring lengths in units of ‘ q -meters.’ From the perspective of another observer $\mathcal{O}$ living outside the fractal, a q -meter is the α -power of an ordinary ‘integer’ meter (or ‘ x -meter,’ or just a ‘meter’). *Vice versa*, the x -rod of an integer observer $\mathcal{O}$ measures meters, but from the perspective of the fractal observer $\mathcal{O}_{\mathcal{F}}$ an x -rod measures $(1/\alpha)$ -powers of q -lengths, not q -lengths. If one did not make this correspondence, one would be insisting in measuring the coast of Britain [43] with an infinitesimal but ordinary integer rod, getting infinity as a result.
- *Multi-scale geometry.* In the multi-fractal case, one replaces $\mathcal{O}_{\mathcal{F}}$ with a microscopic observer measuring the fine scales of a multi-fractal and $\mathcal{O}$ with a macroscopic observer living at scales of the multi-fractal large enough so that the set appears smooth ($d_H \sim D$). Suppose the multi-fractional system is characterized by N charges α_n , corresponding to N asymptotic regimes. For each regime, the geometric coordinates q_n^μ depend on a given α_n . At scales $\ell \sim \ell_1$, corresponding to QEG scales $L(\ell) \sim L_1 = L(\ell_1)$, the QEG momentum is given by eq. (3.5), $p_{\text{QEG}} \sim (p_{\text{frac}})^{\alpha_1} \leftrightarrow q_1^{-1}$; at scales $\ell \sim \ell_2$, $p_{\text{QEG}} \sim (p_{\text{frac}})^{\alpha_2} \leftrightarrow q_2^{-1}$; and so on. Approximately, all the discussion above and below holds replacing q with the effective scale-dependent coordinate q_{eff} , eq. (2.7), so that

$$p_{\text{QEG}}(L) \sim p_{\text{eff}}(\ell) = \frac{(p_{\text{frac}})^{\alpha_{\text{eff}}(\ell)}}{\Gamma[1 + \alpha_{\text{eff}}(\ell)]} \longleftrightarrow q_{\text{eff}}^{-1}(\ell). \quad (3.7)$$

The duality is the first relation $\sim$ of this equation. In this case, the heuristic relation between L and ℓ becomes transcendental, $L \sim \ell^{\alpha_{\text{eff}}(\ell)} / \Gamma[\alpha_{\text{eff}}(\ell) + 1]$.

In QEG, the RG scaling stems from the comparison of any given scale $1/k = L$ with a classical scale $1/k_0 = \ell$. To get finite results, the rod to use must be k -adapted, yet k_0 -dependent. Saying that $k = p_{\text{QEG}}$ is the momentum conjugate to some $q(x)$ is the precise transposition of the idea of getting a k -adapted measurement (in fractional language, a q -rod, suitable for fractals) but dependent on a classical scale in a certain way motivated by the RG flow (i.e., q is a specific function of x given by fractional geometry). Thus, the following correspondences ensue:



Since $L = \ell^\alpha$ and α is the Hausdorff dimension (along one direction), one concludes that *the very concept of dimensional flow is, in fact, the notion of adapted rod*.

Notice that, in fractional theory, the non-trivial measure $\varrho_\alpha(x)$ is not a background but the prescription of the rods by which one makes measurements during the RG flow. At least in the asymptotic regimes of constant spectral dimension, renormalization group and fractal geometry are just two different languages describing a similar geometry (not the same, since symmetries are different; see below). In QEG one does not change the measure but changes the momenta (hence the rods), while in multi-fractional spaces one does not change the momenta but changes the measure (hence the rods again): the two pictures are complementary. Recalling the above analogy of the fractal examined by ‘fractal’ and ‘integer’ observers, one case (fractional theory) corresponds to the integer observer $\mathcal{O}$ measuring the fractal (at any of its scale) with his own x -rods, while the other (QEG) corresponds to the fractal observer $\mathcal{O}_{\mathcal{F}}$ measuring $\mathcal{F}$, at all scales (see table 1).

	Multi-fractional spacetimes		QEG spacetimes
		mapping	
Coordinate	x	$q_{\text{eff}}(x, \ell)$	q
Physical momentum	p_{frac}	$p_{\text{eff}}(\ell)$	p_{QEG}
Scale dependence	implicit	explicit	implicit
Probed scale	$\ell = p_{\text{frac}}^{-1}$	$L = p_{\text{eff}}^{-1}$	$L = p_{\text{QEG}}^{-1}$
Rods adapted via	measure	momenta	momenta
Laplacian	$\mathcal{K}_\alpha \sim \partial_x^2$		$\square^{1/\alpha} \sim \partial_q^{2/\alpha}$

Table 1. QEG vs. multi-fractional picture.

By definition, physical momentum changes according to the scale of the experiment ($p_{\text{QEG}} \sim L^{-1}$, $p_{\text{frac}} \sim \ell^{-1}$), but a scale dependence hidden in the momentum/coordinate

structure is always *implicit*. The mapping to multi-fractional theory makes this scale dependence explicit. To summarize, (a) in QEG the use of q -rods imply that the relation between momenta p_{QEG} and probed scales L is fixed throughout dimensional flow and, as a consequence, the Hausdorff dimension of QEG spacetime does not change, $d_{\text{H}} = D$; (b) in multi-fractional spacetimes, the relation between momenta p_{frac} and probed scales ℓ is also fixed, but the measure changes, which leads to a non-constant Hausdorff dimension $d_{\text{H}} \leq D$; (c) Writing multi-fractional theory with coordinates (2.7) and momenta (3.7) reproduces the QEG situation because the measure is trivialized and scale dependence is absorbed and made *explicit* into the coordinate system.

Lorentz-symmetry violation is manifest in multi-fractional theory⁵ but not usually expected in asymptotic safety. The mapping, however, was done at the level of the action measure between geometric coordinates q in multi-fractional theory and momenta p_{QEG} in QEG. The measure in geometric coordinates is Lorentz invariant, $d^D q$, but the multi-fractional Laplace–Beltrami operator is not Lorentz invariant either in x or in q coordinates, due to the weight factors, and differs from its QEG counterpart (see the last line of table 1, where we made use of eqs. (1.3) and (3.4)). Hence the two theories are physically different although their dimensional flows share a common description.

3.2 Multi-fractional RG flow and Hořava–Lifshitz gravity

To mimick the results of Hořava–Lifshitz (HL) gravity, we look at an anisotropic fractional model with

$$\alpha_0 = 1, \quad \alpha_i = \frac{1}{z} = \frac{1}{D-1}, \quad i = 1, \dots, D-1. \quad (3.8)$$

The spatial fractional charge α_i is nothing but the inverse of the critical exponent z . In particular, in four dimensions $\alpha_i = 1/3$. The coordinates correspondence runs as

$$x_{\text{frac}}^0 = x_{\text{HL}}^0 = t, \quad q^i = x_{\text{HL}}^i. \quad (3.9)$$

The measure then reads $d\varrho_\alpha = d\varrho_{\text{HL}} = dt d^{D-1} \mathbf{q}$. Physical momenta are defined consequently: $p_{\text{frac}}^0 = p_{\text{HL}}^0$, $(p_{\text{frac}}^i)^\alpha \sim p_{\text{HL}}^i$.

To get a multi-scale geometry, one applies perturbative RG considerations to build the total action with a hierarchy of differential Laplacian operators, from order $2z$ (UV) to 2 (IR). While in HL gravity the UV spectral dimension is anomalous due to the higher-order Laplacian, in the fractional mapping it is so because of the non-trivial fractional charges.

Again, symmetries and action differ with respect to multi-fractional theory taken as a stand-alone proposal, so the physics is ultimately inequivalent. In the latter case, the Laplace–Beltrami operator is isotropic, eq. (1.2), $\mathcal{K}_\alpha \sim -p_t^2 + \partial_x^2$, while the operator corresponding to HL spacetime in the UV is $\mathcal{K} \sim -\partial_t^2 + \partial_q^{2/\alpha}$. This is the reason why, in particular, in $D = 2 + 1$ dimensions the two models predict a different UV spectral dimension, $d_S \sim 3/2$ (as in QEG) in the fractional case and $d_S = 2$ in the HL case. Our purpose

⁵If one attempted to work with a measure preserving as many Lorentz symmetries as possible, for instance rotations ($v_\alpha = |\mathbf{x}|^{\alpha-1}$), one would obtain a model with the correct anomalous scaling but which is technically untractable after a few steps [4–6]. There is no obvious (and perhaps even non-obvious) momentum transform for non-factorizable measures, the propagator of fields cannot be computed rigorously, and so on. The factorized form (2.1b) is crucial.

here was to show how the main geometric features of HL dimensional flow can be easily reproduced by the multi-fractional framework.

4 Discussion

A twofold purpose of multi-fractional geometry is, when regarded as an independent theory, to provide a general analytic explanation of dimensional flow [12] and, when regarded as an effective model, to reproduce the flow in other theories and give a complementary description. Both aims can be stated and reached in such a generality because multi-fractional spacetimes are constructed with the wide-scope tools of multi-fractal geometry, anomalous transport and complex systems. We have applied this *passe-partout* spirit to the multi-scale behaviour of QEG and Hořava–Lifshitz effective spacetimes. In particular, we have revisited their RG flow within fractal geometry.

In general, the multi-scale nature of the renormalization group is akin to the structure found in multi-fractal systems.⁶ At the level of the Laplacian operator, the correspondence between different theories is based on the following heuristic observation. In the UV, the spectral dimension of multi-fractional, QEG and HL spacetimes is 2 in four dimensions. Calling γ_μ half the order of the Laplacian along the direction μ and allowing for a non-trivial position-space measure, the spectral dimension for a non-fractional diffusion operator is $d_S = \sum_\mu \alpha_\mu / \gamma_\mu$ [12]. This formula is invariant under the interchange

$$\alpha_\mu \longleftrightarrow \frac{1}{\gamma_\mu}, \quad (4.1)$$

mapping theories with $(\alpha_\mu, \gamma_\mu) = (1, \gamma_\mu)$ (ordinary measure, higher-order Laplacians) to a fractional spacetime with $(\alpha_\mu, \gamma_\mu) = (1/\gamma_\mu, 1)$ (non-trivial measure, second-order Laplacian). This is realized by eq. (3.4) in asymptotic safety and by eq. (3.8) in HL gravity. Despite this intriguing hint, when one looks at the details of each Laplacian one finds differences in the symmetries. Although these are intrinsic and cannot be bypassed, it is nonetheless desirable to clarify the correspondence in the dimensional flow and recast it without invoking the form of the Laplacian. Here we did so and pointed out how a more precise quantitative comparison between quantum field theory renormalization and fractal geometry is based upon the identification of coordinates, measure in position space, and physical momenta.

As shown in section 3.1, the natural coordinates in QEG do not correspond to the x variables in fractional spacetime but to the ‘geometric’ coordinates q . This implies that the physical momentum of QEG is the same as in non-commutative geometries with fractal properties, although non-commutative space is spanned by $\{x\}$, not by $\{q\}$, and with a non-commutative product rule [15]. Thus, the physical picture of these non-commutative geometries is somewhat hybrid between quantum gravity and fractional spacetimes. Interestingly, this observation pinpoints one of the key differences between all these independent approaches to quantum geometry: namely, *the choice of momentum space*. In other words,

⁶Whether there is a deeper link between fractals and anomalous diffusion on one hand and the hidden tree structure of Feynman graphs ([44–46] and references therein) on the other hand is a matter of speculation.

what distinguishes one model from another is, apart from the symmetries, the scale identification. This is also the leverage point where one can start drawing a quantitative mapping between these theories, as shown here and in [15]. A better understanding of these relations may be a promising line of research for the near future.

Acknowledgments

The author thanks A. Eichhorn, V. Rivasseau, F. Saueressig and J. Thürigen for useful discussions. This work is supported by a Ramón y Cajal tenure-track fellowship.

References

- [1] G. 't Hooft, *Dimensional reduction in quantum gravity*, in *Salamfestschrift*, A. Ali, J. Ellis and S. Randjbar-Daemi (eds.), World Scientific, Singapore (1993) [[gr-qc/9310026](#)].
- [2] S. Carlip, *Spontaneous dimensional reduction in short-distance quantum gravity?*, in *Proceedings of the XXV Max Born Symposium*, Wrocław, Poland, 2009 (to be published) [[arXiv:0909.3329](#)].
- [3] S. Carlip, *The small scale structure of spacetime*, in *Foundations of Space and Time*, G. Ellis, J. Murugan and A. Weltman (eds.), Cambridge University Press, Cambridge U.K. (2012) [[arXiv:1009.1136](#)].
- [4] G. Calcagni, *Fractal universe and quantum gravity*, *Phys. Rev. Lett.* **104** (2010) 251301 [[arXiv:0912.3142](#)].
- [5] G. Calcagni, *Quantum field theory, gravity and cosmology in a fractal universe*, *JHEP* **1003** (2010) 120 [[arXiv:1001.0571](#)].
- [6] G. Calcagni, *Gravity on a multifractal*, *Phys. Lett. B* **697** (2011) 251 [[arXiv:1012.1244](#)].
- [7] G. Calcagni, *Discrete to continuum transition in multifractal spacetimes*, *Phys. Rev. D* **84** (2011) 061501(R) [[arXiv:1106.0295](#)].
- [8] G. Calcagni, *Geometry of fractional spaces*, *Adv. Theor. Math. Phys.* **16** to appear [[arXiv:1106.5787](#)].
- [9] G. Calcagni, *Geometry and field theory in multi-fractional spacetime*, *JHEP* **1201** (2012) 065 [[arXiv:1107.5041](#)].
- [10] G. Calcagni and G. Nardelli, *Momentum transforms and Laplacians in fractional spaces*, [arXiv:1202.5383](#).
- [11] G. Calcagni, *Diffusion in quantum geometry*, *Phys. Rev. D* **86** (2012) 044021 [[arXiv:1204.2550](#)].
- [12] G. Calcagni, *Diffusion in multi-fractional spacetimes*, [arXiv:1205.5046](#).
- [13] G. Calcagni, G. Nardelli and M. Scalisi, *Quantum mechanics in fractional and other anomalous spacetimes*, *J. Math. Phys.* (to be published) [[arXiv:1207.4473](#)].
- [14] G. Calcagni, *Introduction to multifractional spacetimes*, to appear in Proceedings of the Sixth International School on Field Theory and Gravitation, Petrópolis, Brazil, 23-27 April 2012 [[arXiv:1209.1110](#)].

- [15] M. Arzano, G. Calcagni, D. Oriti and M. Scalisi, *Fractional and noncommutative spacetimes*, *Phys. Rev. D* **84** (2011) 125002 [[arXiv:1107.5308](#)].
- [16] S. Weinberg, *Ultraviolet divergences in quantum gravity*, in *General relativity, an Einstein centenary survey*, S.W. Hawking and W. Israel eds., Cambridge University Press, Cambridge U.K. (1979).
- [17] M. Reuter, *Nonperturbative evolution equation for quantum gravity*, *Phys. Rev. D* **57** (1998) 971 [[hep-th/9605030](#)].
- [18] M. Niedermaier, *The asymptotic safety scenario in quantum gravity: an introduction*, *Class. Quantum Grav.* **24** (2007) R171 [[gr-qc/0610018](#)].
- [19] D.F. Litim, *Renormalisation group and the Planck scale*, *Phil. Trans. Roy. Soc. Lond. A* **369** (2011) 2759 [[arXiv:1102.4624](#)].
- [20] M. Reuter and F. Saueressig, *Fractal space-times under the microscope: a renormalization group view on Monte Carlo data*, *JHEP* **1112** (2011) 012 [[arXiv:1110.5224](#)].
- [21] M. Reuter and F. Saueressig, *Asymptotic safety, fractals, and cosmology*, in *Quantum Gravity and Quantum Cosmology*, G. Calcagni, L. Papantonopoulos, G. Siopsis and N. Tsamis (eds.), Springer-Verlag, Berlin Germany (2012, to be published) [[arXiv:1205.5431](#)].
- [22] G. Calcagni, A. Eichhorn and F. Saueressig, in progress.
- [23] P. Hořava, *Quantum gravity at a Lifshitz point*, *Phys. Rev. D* **79** (2009) 084008 [[arXiv:0901.3775](#)].
- [24] P. Hořava, *Spectral dimension of the universe in quantum gravity at a Lifshitz point*, *Phys. Rev. Lett.* **102** (2009) 161301 [[arXiv:0902.3657](#)].
- [25] G. Calcagni and G. Nardelli, *Symmetries and propagator in multi-fractional scalar field theory*, in progress.
- [26] D. Harte, *Multifractals: theory and applications*, Chapman & Hall/CRC, Boca Raton U.S.A. (2001).
- [27] D. Hochberg and J.T. Wheeler, *Spacetime dimension from a variational principle*, *Phys. Rev. D* **43** (1991) 2617.
- [28] L. Nottale, *Scale relativity and fractal space-time: theory and applications*, *Found. Sci.* **15** (2010) 101 [[arXiv:0812.3857](#)].
- [29] C. Bagnuls and C. Bervillier, *Exact renormalization group equations: an introductory review*, *Phys. Rept.* **348** (2001) 91 [[hep-th/0002034](#)].
- [30] J. Berges, N. Tetradis and C. Wetterich, *Non-perturbative renormalization flow in quantum field theory and statistical physics*, *Phys. Rept.* **363** (2002) 223 [[hep-ph/0005122](#)].
- [31] J. Polonyi, *Lectures on the functional renormalization group method*, *Central Eur. J. Phys.* **1** (2003) 1 [[hep-th/0110026](#)].
- [32] O.J. Rosten, *Fundamentals of the exact renormalization group*, *Phys. Rept.* **511** (2012) 177 [[arXiv:1003.1366](#)].
- [33] O. Lauscher and M. Reuter, *Fractal spacetime structure in asymptotically safe gravity*, *JHEP* **0510** (2005) 050 [[hep-th/0508202](#)].
- [34] S. Rechenberger and F. Saueressig, *R^2 phase-diagram of QEG and its spectral dimension*, *Phys. Rev. D* **86** (2012) 024018 [[arXiv:1206.0657](#)].

- [35] E.M. Lifshitz, *Zh. Eksp. Teor. Fiz.* **11** (1941) 255.
- [36] E.M. Lifshitz, *Zh. Eksp. Teor. Fiz.* **11** (1941) 269.
- [37] E. Ardonne, P. Fendley and E. Fradkin, *Topological order and conformal quantum critical points*, *Annals Phys.* **310** (2004) 493 [[cond-mat/0311466](#)].
- [38] J. Cardy, *Scaling and renormalization in statistical physics*, Cambridge University Press, Cambridge U.K. (1996).
- [39] P.M. Chaikin and T.C. Lubensky, *Principles of condensed matter physics*, Cambridge University Press, Cambridge U.K. (1995).
- [40] P. Hořava, *General covariance in gravity at a Lifshitz point*, *Class. Quantum Grav.* **28** (2011) 114012 [[arXiv:1101.1081](#)].
- [41] T.P. Sotiriou, M. Visser and S. Weinfurtner, *Spectral dimension as a probe of the ultraviolet continuum regime of causal dynamical triangulations*, *Phys. Rev. Lett.* **107** (2011) 131303 [[arXiv:1105.5646](#)].
- [42] R. Iengo, J.G. Russo and M. Serone, *Renormalization group in Lifshitz-type theories*, *JHEP* **0911** (2009) 020 [[arXiv:0906.3477](#)].
- [43] B. Mandelbrot, *How long is the coast of Britain? Statistical self-similarity and fractional dimension*, *Science* **156** (1967) 636.
- [44] W. Zimmermann, *Convergence of Bogoliubov's method for renormalization in momentum space*, *Commun. Math. Phys.* **15** (1969) 208.
- [45] D. Kreimer, *Combinatorics of (perturbative) quantum field theory*, *Phys. Rept.* **363** (2002) 387 [[hep-th/0010059](#)].
- [46] G. Gallavotti, *Perturbation theory*, [arXiv:0711.2544](#).